

DESCRIPTION

AQUEOUS COMPOSITION COMPRISING THIAZOLE DERIVATIVE

TECHNICAL FIELD


The present invention relates to an aqueous
5 composition comprising a specific thiazole derivative.

BACKGROUND ART

Vascular adhesion protein-1 (hereinafter to be
abbreviated as VAP-1) is an amine oxidase (semicarbazide
sensitive amine oxidase, SSAO) which is abundant in human
10 plasma, and shows remarkably increased expression in
vascular endothelium and vascular smooth muscle of the
inflammatory region. While the physiological role of VAP-1
has not been clarified until recently, VAP-1 gene was
cloned in 1998, and VAP-1 has been reported to be a
15 membrane protein that regulates rolling and migration of
lymphocyte and NK cell as an adhesion molecule under-
regulation of expression by inflammatory cytokine.
Although the amine to be a substrate is unknown, it is
considered to be methylamine generated in any part of
20 living organisms. It is also known that hydrogen peroxide
and aldehydes produced due to the amine oxidase activity in
the molecule are important factors of adhesion activity.

Thiazole derivatives represented by formula (A)
below are useful as VAP-1 inhibitor (US Patent Publication
25 No. 20040259923A1 published on December 23, 2004, the

contents of which is incorporated by reference).




wherein

R^1 is acyl;

5 X is a bivalent residue derived from optionally substituted thiazole;

Y is a bond, lower alkylene, lower alkenylene or -CONH-; and

Z is a group of the formula:




wherein R^2 is a group of the formula: -A-B-D-E

wherein A is a bond, lower alkylene, -NH- or -SO₂-;

B is a bond, lower alkylene, -CO- or -O-;

D is a bond, lower alkylene, -NH- or -CH₂NH-; and

15 E is optionally protected amino, -N=CH₂,



wherein

Q is -S- or -NH-; and

R^3 is hydrogen, lower alkyl, lower alkylthio or

20 -NH- R^4 wherein R^4 is hydrogen, -NH₂ or

lower alkyl;

or a pharmaceutically acceptable salt thereof.


Usually, in preparing an aqueous dosage forms such as injection, eye drops and the like, sodium chloride is added to adjust osmotic pressure of the solution. However, in preparing an aqueous composition comprising the
5 above thiazole derivative, there is a problem that the presence of sodium chloride in the solution decreases the solubility of the above thiazole derivative and precipitates the derivative.

SUMMARY OF THE INVENTION

10 An object of the present invention is to provide an aqueous composition comprising a certain thiazole derivative which is clear and stable, and can be stored for a long time.

Thus, the present invention provides

15 [1] An aqueous composition comprising a compound of formula (I) [hereinafter sometimes referred to as Compound (I)]:




wherein

20 R^1 is acyl;

X is a bivalent residue derived from optionally substituted thiazole;

Y is a bond, lower alkylene, lower alkenylene or -CONH-; and

25 Z is a group of the formula:




wherein R² is a group of the formula: -A-B-D-E

wherein A is a bond, lower alkylene, -NH- or -SO₂-;

B is a bond, lower alkylene, -CO- or -O-;

5 D is a bond, lower alkylene, -NH- or -CH₂NH-, provided that when B is -CO- or -O-, D is not a bond; and

E is optionally protected amino, -N=CH₂,



wherein Q is -S- or -NH-; and


10 R³ is hydrogen, lower alkyl, lower alkylthio or -NH-R⁴

wherein R⁴ is hydrogen, -NH₂ or lower alkyl;


or a pharmaceutically acceptable salt thereof,

and an additive selected from the group consisting of
15 polyol, sugar, sugar alcohol, boric acid or its salt,
and water.


[2] The composition of [1], wherein Z of the compound (I) is a group of the formula:



20 wherein R² is a group of the formula:




(wherein G is a bond, $\text{-NHCOCH}_2\text{-}$ or lower alkylene and R^4 is hydrogen, -NH_2 or lower alkyl); -NH_2 ; $\text{-CH}_2\text{NH}_2$; $\text{-CH}_2\text{ONH}_2$; $\text{-CH}_2\text{ON=CH}_2$;




5 or a pharmaceutically acceptable salt thereof.

[3] The composition of [2], wherein R^2 of the compound (I) is a group of the formula:



10 (wherein G is a bond, $\text{-NHCOCH}_2\text{-}$ or lower alkylene and R^4 is hydrogen or lower alkyl); $\text{-CH}_2\text{NH}_2$; $\text{-CH}_2\text{ONH}_2$; $\text{-CH}_2\text{ON=CH}_2$;



or a pharmaceutically acceptable salt thereof.

15 [4] The composition of any of [1] to [3], wherein R^1 of the compound (I) is alkylcarbonyl and X is a bivalent residue derived from thiazole optionally substituted by methylsulfonylbenzyl, or a pharmaceutically acceptable salt thereof.

20 [5] The composition of [1], wherein the compound (I) is N-{4-[2-(4-{[amino(imino)methyl]amino}phenyl)ethyl]-1,3-thiazol-2-yl}acetamide,

N-{4-[2-(4-{[amino(imino)methyl]amino}phenyl)ethyl]-5-[4-(methylsulfonyl)benzyl]-1,3-thiazol-2-yl]acetamide,

N-{4-[2-(4-{[hydrazino(imino)methyl]amino}phenyl)ethyl]-5-[4-(methylsulfonyl)benzyl]-1,3-thiazol-2-yl]acetamide,

5 N-{4-[2-(4-{[hydrazino(imino)methyl]amino}phenyl)ethyl]-1,3-thiazol-2-yl]acetamide, or

N-(4-{2-[4-(2-{[amino(imino)methyl]amino}ethyl)phenyl]ethyl}-1,3-thiazol-2-yl)acetamide,

or a pharmaceutically acceptable salt thereof.

10 DETAILED DESCRIPTION OF THE INVENTION

In the above and subsequent descriptions of the present specification, suitable examples and illustration of the various definitions to be included within the scope of the invention are explained in detail as follows.

15 Suitable "halogen" includes fluorine, chlorine, bromine and iodine.

The term "lower" is used to intend a group having 1 to 6, preferably 1 to 4, carbon atom(s), unless otherwise provided.

20 Suitable "lower alkyl" includes straight or branched alkyl having 1 to 6 carbon atom(s), such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, tert-pentyl and hexyl, in which more preferred one is C₁-C₄ alkyl.

25 Suitable "lower alkylthio" includes lower alkylthio

containing the above lower alkyl, such as methylthio, ethylthio, propylthio, isopropylthio, butylthio, isobutylthio, sec-butylthio, tert-butylthio, pentylthio, tert-pentylthio and hexylthio.

5 Suitable "lower alkylene" includes straight or branched alkylene having 1 to 6 carbon atom(s), such as methylene, ethylene, trimethylene, tetramethylene, propylene, ethylidene and propylidene, in which more preferred one is C₁-C₄ alkylene.

10 Suitable "lower alkenylene" includes straight or branched alkenylene having 2 to 6 carbon atom(s), such as

-CH=CH-, -CH₂-CH=CH-, -CH₂-CH=CH-CH₂-, -CH₂-CH₂-CH=CH-,

-CH=CH-CH=CH-, -CH=CH-CH₂-CH₂-CH₂-, -CH=CH-CH=CH-CH₂-CH₂-

and

15 -CH=CH-CH=CH-CH=CH-, in which more preferred one is C₂-C₄ alkenylene.

The above lower alkenylene may be in E or Z form, respectively. Thus, those skilled in the art will recognize that the lower alkenylene includes all E, Z-
20 structures when it has 2 or more double bonds.

Suitable "aryl" includes C₆-C₁₀ aryl such as phenyl and naphthyl, in which more preferred one is phenyl. The "aryl" may be substituted by 1 to 3 substituent(s) and the substitution sites are not particularly limited.

25 Suitable "aralkyl" includes aralkyl wherein the aryl

moiety has 6 to 10 carbon atoms [i.e. the aryl moiety is C₆-C₁₀ aryl of the above "aryl"] and the alkyl moiety has 1 to 6 carbon atom(s) [i.e. the alkyl moiety is C₁-C₆ alkyl of the above "lower alkyl"], such as benzyl, phenethyl, 1-naphthylmethyl, 2-naphthylmethyl, 3-phenylpropyl, 4-phenylbutyl and 5-phenylpentyl.

The "optionally protected amino" means that an amino group may be protected with a suitable protecting group according to a method known *per se*, such as the methods described in Protective Groups in Organic Synthesis, published by John Wiley and Sons (1980), and the like. The suitable "protecting group" includes tert-butoxycarbonyl (i.e., Boc), an acyl group as mentioned below, substituted or unsubstituted aryl(lower)alkylidene [e.g., benzylidene, hydroxybenzylidene, etc.], aryl(lower)alkyl such as mono-, di- or triphenyl-(lower)alkyl [e.g., benzyl, phenethyl, benzhydryl, trityl, etc.] and the like.

Suitable "optionally protected amino" includes amino and tert-butoxycarbonylamino (i.e. -NHBoc).

Suitable "heterocycle" includes "aromatic heterocycle" and "non-aromatic heterocycle".

Suitable "aromatic heterocycle" includes 5 to 10-membered aromatic heterocycle containing 1 to 3 heteroatom(s) selected from nitrogen, oxygen and sulfur atoms besides carbon atom(s), and includes, for example,

thiophene, furan, pyrrole, imidazole, pyrazole, thiazole, isothiazole, oxazole, isoxazole, pyridine, pyridazine, pyrimidine, pyrazine and the like.

Suitable "non-aromatic heterocycle" includes 5 to 10-membered non-aromatic heterocycle containing 1 to 3 heteroatom(s) selected from nitrogen, oxygen and sulfur atoms besides carbon atom(s), and includes, for example, pyrrolidine, imidazoline, pyrazolidine, pyrazoline, piperidine, piperazine, morpholine, thiomorpholine, dioxolan, oxazolidine, thiazolidine, triazolidine and the like.

Suitable "acyl" includes acyl having 1 to 20 carbon atom(s), such as formyl, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl and aralkyloxycarbonyl.

Suitable "alkylcarbonyl" includes alkylcarbonyl wherein the alkyl moiety has 1 to 6 carbon atom(s) [i.e. the alkyl moiety is C₁-C₆ alkyl of the above "lower alkyl"], such as acetyl, propionyl, butyryl, isobutyryl, valeryl, isovaleryl, pivaloyl, hexanoyl and heptanoyl, in which more preferred one is C₁-C₄ alkyl-carbonyl.


Suitable "arylcarbonyl" includes arylcarbonyl wherein the aryl moiety has 6 to 10 carbon atom(s) [i.e. the aryl moiety is C₆-C₁₀ aryl of the above "aryl"], such as benzoyl and naphthoyl.

Suitable "alkoxycarbonyl" includes alkoxycarbonyl

wherein the alkoxy moiety has 1 to 6 carbon atom(s), such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, sec-butoxycarbonyl, tert-butoxycarbonyl, pentyloxycarbonyl, tert-pentyloxycarbonyl and hexyloxycarbonyl, in which more preferred one is alkoxycarbonyl wherein the alkoxy moiety has 1 to 4 carbon atom(s).

Suitable "aralkyloxycarbonyl" includes aralkyloxycarbonyl wherein the aryl moiety has 6 to 10 carbon atom(s) [i.e. the aryl moiety is C₆-C₁₀ aryl of the above "aryl"] and the alkyl moiety has 1 to 6 carbon atom(s) [i.e. the alkyl moiety is C₁-C₆ alkyl of the above "lower alkyl"], such as benzyloxycarbonyl, phenethyloxycarbonyl, 1-naphthylmethyloxycarbonyl, 2-naphthylmethyloxycarbonyl, 3-phenylpropyloxycarbonyl, 4-phenylbutyloxycarbonyl and 5-phenylpentyloxycarbonyl.

Suitable "bivalent residue derived from thiazole" of the "bivalent residue derived from optionally substituted thiazole" includes



The "thiazole" may have 1 to 3 substituent(s) and the substitution sites are not particularly limited.

Suitable "substituent" of the above "optionally substituted thiazole" includes, for example,

(1) halogen which is as defined above;

(2) alkoxy carbonyl which is as defined above, such as ethoxy carbonyl;

(3) optionally substituted aryl, which aryl is as defined above and the substitution sites are not particularly limited, such as phenyl and 4-(methylsulfonyl)phenyl;

(4) a group of the formula: $-\text{CONR}^a\text{R}^b$ wherein R^a is hydrogen, lower alkyl, aryl or aralkyl and R^b is hydrogen, lower alkyl, aryl or aralkyl, wherein the lower alkyl, aryl and aralkyl are as defined above, such as N-methylaminocarbonyl, N-phenylaminocarbonyl, N,N-dimethylaminocarbonyl and N-benzylaminocarbonyl;

(5) a group of the formula: $-\text{CONH}-(\text{CH}_2)_k\text{-aryl}$ wherein k is an integer of 0 to 6; the aryl is as defined above, which may have 1 to 5 substituent(s) selected from the group consisting of $-\text{NO}_2$, $-\text{SO}_2\text{-(lower alkyl)}$ wherein the lower alkyl is as defined above, $-\text{CF}_3$ and $-\text{O-aryl}$ wherein the aryl is as defined above, and the substitution sites are not particularly limited;

(6) a group of the formula: $-\text{CONH}-(\text{CH}_2)_m\text{-heterocycle}$ wherein m is an integer of 0 to 6; the heterocycle is as defined above, such as pyridine;

(7) a group of the formula: $-\text{CO-heterocycle}$ wherein the heterocycle is as defined above, such as

pyrrolidine, piperidine, piperazine, thiomorpholine, which may have 1 to 5 substituent(s) selected from the group consisting of -CO-(lower alkyl) wherein the lower alkyl is as defined above, -CO-O-(lower alkyl) wherein the lower alkyl is as defined above, -SO₂-(lower alkyl) wherein the lower alkyl is as defined above, oxo (i.e. =O) and a group of the formula: -CONR^cR^d wherein R^c is hydrogen, lower alkyl, aryl or aralkyl and R^d is hydrogen, lower alkyl, aryl or aralkyl wherein the lower alkyl, aryl and aralkyl are as defined above, and the substitution sites are not particularly limited;

(8) a group of the formula: -(CH₂)_n-aryl wherein n is an integer of 1 to 6; the aryl is as defined above, which may have 1 to 5 substituent(s) selected from the group consisting of -S-(lower alkyl) wherein the lower alkyl is as defined above, -SO₂-(lower alkyl) wherein the lower alkyl is as defined above, -CO₂-(lower alkyl) wherein the lower alkyl is as defined above, -NHCO-O-(lower alkyl) wherein the lower alkyl is as defined above and a group of the formula: -CONR^eR^f wherein R^e is hydrogen, lower alkyl, aryl or aralkyl and R^f is hydrogen, lower alkyl, aryl or aralkyl wherein the lower alkyl, aryl and aralkyl are as defined above, and the substitution sites are not particularly limited;

(9) a group of the formula: -(CH₂)_o-heterocycle


wherein o is an integer of 0 to 6; the heterocycle is as defined above, such as pyrrolidine, piperidine, piperazine, morpholine, thiomorpholine, which may have 1 to 5 substituent(s) selected from the group consisting of oxo (i.e. =O); -CO-(lower alkyl) wherein the lower alkyl is as defined above; -CO-O-(lower alkyl) wherein the lower alkyl is as defined above; -SO₂-(lower alkyl) wherein the lower alkyl is as defined above; -CO-(heterocycle) wherein the heterocycle is as defined above such as pyrrolidine, piperazine and morpholine, which may have 1 to 5 substituent(s) selected from the group consisting of lower alkyl and halogen, wherein the lower alkyl and halogen are as defined above, and the substitution sites are not particularly limited; and a group of the formula: -CONR^gR^h wherein R^g is hydrogen, lower alkyl, aryl or aralkyl and R^h is hydrogen, lower alkyl, aryl or aralkyl wherein the lower alkyl, aryl and aralkyl are as defined above, and the substitution sites are not particularly limited;

(10) a group of the formula: -(CH₂)_p-NRⁱR^j

wherein p is an integer of 0 to 6; Rⁱ is hydrogen, acyl, lower alkyl, aryl or aralkyl and R^j is hydrogen, acyl, lower alkyl, aryl or aralkyl wherein the acyl, lower alkyl, aryl and aralkyl are as defined above, and the lower alkyl may have 1 to 5 substituent(s) selected from the group consisting of a group of the formula: -CONR^kR^l wherein R^k is

hydrogen, lower alkyl, aryl or aralkyl and R^1 is hydrogen, lower alkyl, aryl or aralkyl wherein the lower alkyl, aryl and aralkyl are as defined above, and the substitution sites are not particularly limited;

5 (11) a group of the formula:



wherein q is an integer of 0 to 6; the lower alkyl is as defined above; R^m is hydrogen, aralkyl which is as defined above, or alkyl which is as defined above, which may be substituted by 1 to 3 substituent(s) selected from the group consisting of $-\text{OH}$ and $-\text{CONH}_2$ and the substitution sites are not particularly limited; and T is hydrogen; a group of the formula: $-\text{CONR}^n\text{R}^o$ wherein R^n is hydrogen, lower alkyl, aryl or aralkyl and R^o is hydrogen, lower alkyl, aryl or aralkyl wherein the lower alkyl, aryl and aralkyl are as defined above; $-\text{NH}-\text{CO}-\text{R}^p$ wherein R^p is lower alkyl which is as defined above or aralkyl which is as defined above;

15 $-\text{NH}-\text{SO}_2-(\text{lower alkyl})$ wherein the lower alkyl is as defined above; $-\text{SO}_2-(\text{lower alkyl})$ wherein the lower alkyl is as defined above; $-\text{heterocycle}$ wherein the heterocycle is as defined above, such as pyridine, pyrrolidine and morpholine, which may have 1 to 3 substituent(s) such as oxo (i.e. $=\text{O}$), and the substitution sites are not particularly limited; or
20 $-\text{CO}-(\text{heterocycle})$ wherein the heterocycle is as defined

above, such as piperidine and morpholine; and


(12) a group of the formula: $-(CH_2)_r-CO-NR^tR^u$


wherein r is an integer of 1 to 6; R^t is hydrogen, lower alkyl, aryl or aralkyl and R^u is hydrogen, lower alkyl, aryl or aralkyl wherein the lower alkyl, aryl and aralkyl are as defined above.

The substitution site on the aryl or heterocycle is any suitable position thereof, but not particularly limited. Preferable "substituent" of the above "optionally substituted thiazole" is methylsulfonylbenzyl.

The substitution sites of R^2 on the phenyl in Compound (I) is not particularly limited.

When Z is a group of the formula:



the substitution sites on the group are not particularly limited.  is particularly preferable.

Any nitrogen atom in the amino (i.e. $-NH_2$), imino (i.e. $=NH$ or $-NH-$) or the like contained in Compound (I) may be protected according to the methods, which are known to those skilled in the art, such as the methods described in Protective Groups in Organic Synthesis, published by John Wiley and Sons (1980), and the like.

When Compound (I) has an asymmetric carbon atom in the structure, those skilled in the art will recognize that

Compound (I) includes all stereoisomers.

Of the above-mentioned compounds, preferred are Compound (I), more preferably,

N-{4-[2-(4-{[amino(imino)methyl]amino}phenyl)ethyl]-1,3-thiazol-2-yl}acetamide (see Structure 1),

N-{4-[2-(4-{[amino(imino)methyl]amino}phenyl)ethyl]-5-[4-(methylsulfonyl)benzyl]-1,3-thiazol-2-yl}acetamide (see Structure 46),

N-{4-[2-(4-{[hydrazino(imino)methyl]amino}phenyl)ethyl]-5-[4-(methylsulfonyl)benzyl]-1,3-thiazol-2-yl}acetamide (see Structure 48),

N-{4-[2-(4-{[hydrazino(imino)methyl]amino}phenyl)ethyl]-1,3-thiazol-2-yl}acetamide (see Structure 56), and

N-(4-{2-[4-(2-{[amino(imino)methyl]amino}ethyl)phenyl]ethyl}-1,3-thiazol-2-yl)acetamide (see Structure 107), particularly N-{4-[2-(4-{[amino(imino)methyl]amino}phenyl)ethyl]-1,3-thiazol-2-yl}acetamide and derivatives thereof.

The term "derivative" is intended to include all compounds derived from the original compound.

The pharmaceutically acceptable salt of Compound (I) of the present invention is nontoxic and a pharmaceutically acceptable conventional salt, which is exemplified by salts with inorganic or organic base such as alkali metal salt (e.g., sodium salt, potassium salt and the like), alkaline

earth metal salt (e.g., calcium salt, magnesium salt and the like), ammonium salt, and amine salt (e.g., triethylamine salt, N-benzyl-N-methylamine salt and the like).

5 The Compound (I) can be also formulated as a pharmaceutically acceptable acid addition salt. Examples of the pharmaceutically acceptable acid addition salts for use in the pharmaceutical composition include those derived from mineral acids, such as hydrochloric, hydrobromic, 10 hydriodic, phosphoric, metaphosphoric, nitric and sulfuric acids, and organic acids, such as tartaric, acetic, citric, malic, lactic, fumaric, benzoic, glycolic, gluconic, succinic and arylsulfonic acids, for example, p-toluenesulfonic acid.

15 As a pharmaceutically acceptable salt of Compound (I) represented by the formula (I), a pharmaceutically acceptable acid addition salt such as (mono-, di- or tri-)hydrochloride and hydriodide, particularly hydrochloride, is preferable.

20 The above-mentioned Compound (I) may be commercially available or can be produced based on a known reference. The composition can be administered in accordance with the present inventive method by any suitable route. Suitable routes of administration include systemic, such as orally 25 or by injection, topical, periocular (e.g., subTenon's),

subconjunctival, intraocular, intravitreal, intracameral, subretinal, suprachoroidal and retrobulbar administrations. The manner in which the VAP-1 inhibitor is administered is dependent, in part, upon whether the treatment of a VAP-1 associated disease is prophylactic or therapeutic.

The composition is preferably administered as soon as possible after it has been determined that a subject such as mammal, specifically a human, is at risk for a VAP-1 associated disease (prophylactic treatments) or has begun to develop a VAP-1 associated disease (therapeutic treatments). Treatment will depend, in part, upon the particular VAP-1 inhibitor to be used, the amount of the VAP-1 inhibitor to be administered, the route of administration, and the cause and extent, if any, of a VAP-1 associated disease realized.

One skilled in the art will appreciate that suitable methods of administering a VAP-1 inhibitor, which is useful in the present inventive method, are available. Although more than one route can be used to administer a particular VAP-1 inhibitor, a particular route can provide a more immediate and more effective reaction than another route. Accordingly, the described routes of administration are merely exemplary and are in no way limiting.

The dose of the composition administered to the administration subject such as animal including human,

particularly a human, in accordance with the present invention should be sufficient to effectuate the desired response in the subject over a reasonable time frame. One skilled in the art will recognize that dosage will depend
5 upon a variety of factors, including the strength of the particular VAP-1 inhibitor to be employed, the age, species, conditions or disease states, and body weight of the subject, as well as the degree of a VAP-1 associated disease. The size of the dose also will be determined by
10 the route, timing and frequency of administration as well as the existence, nature, and extent of any adverse side effects that might accompany the administration of a particular VAP-1 inhibitor and the desired physiological effect. It will be appreciated by one of ordinary skill in
15 the art that various conditions or disease states may require prolonged treatment involving multiple administrations.

Suitable doses and dosage regimens can be determined by conventional range-finding techniques known to those of
20 ordinary skill in the art. Generally, treatment is initiated with smaller dosages, which are less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the optimum effect under the circumstances is reached.

25 Generally, the compound (I) can be administered in the

dose of from about 1 $\mu\text{g/kg/day}$ to about 300 mg/kg/day , preferably from about 0.1 mg/kg/day to about 10 mg/kg/day , which is given in a single dose or 2 to 4 doses a day or in a sustained manner.

5 In the present specification and claims, the "aqueous composition" means clear aqueous solution. The aqueous composition of the present invention may be provided as ophthalmic solution, nasal solution, ear solution, inhalant solution, spray, oral solution, injectable solution for
10 intravenous, intraarterial, subcutaneous, intramuscular, interperitoneal or intraocular administration. Since the additives comprised in the aqueous composition of the present invention, i.e. additives selected from the group consisting of polyol, sugar, sugar alcohol, boric acid or
15 its salt will not affect on solubility of Compound (1), a stable aqueous dosage form with a long shelf life can be provided.

 According to the invention, the additives are those known to be useful for adjusting osmotic pressure of
20 aqueous solution. Examples of the additives used in the present invention may include polyols such as glycerin, polyethylene glycol, propylene glycols and polyvinyl alcohol; saccharides or sugar alcohols such as glucose, sorbitol, mannitol and xylitol; boric acid and its salt.
25 Those additives may be used either solely or in combination

two or more of them depending on the desired formula. Especially preferable additives include glycerin, mannitol, boric acid or its salt.

The concentration of the additives in the aqueous composition of the present invention may be determined so that the osmotic pressure of the composition is adjusted appropriately. The art can determine the concentration based on the kind of the thiazol derivative, the amount of the thiazole derivative as well as the kind and molecular weight of the additives. Typically, the amount of the additive may be 0.001-10w/v%, preferably 0.01-5 w/v% based on total volume of the composition.

The aqueous composition of the present invention may further comprises other additives which are generally used in manufacturing medical compositions, such as a buffering agent, a preservative, a stabilizer and a thickening-agent so long as those additive affect on the solubility of the thiazole derivative or Compound 1 of the present invention. Examples of said additives may include preservatives such as benzalkonium chloride, chlorobutanol and paraoxybenzoates; thickening agent such as povidone and methylcellulose.

The aqueous composition of the present invention may further comprise or may be co-administrated with a pharmaceutically active compound other than Compound (I).

By "co-administration" is meant administration before, concurrently with, e.g., in combination with the VAP-1 inhibitor or Compound (I) thiazole derivative in the same formulation or in separate formulations, or after administration of the VAP-1 inhibitor as described above.











For example, corticosteroids, prednisone, methylprednisolone, dexamethasone, or triamcinolone acetonide, or noncorticosteroid anti-inflammatory compounds, such as ibuprofen or flubiprofen, can be co-administered.











Similarly, vitamins and minerals, e.g., zinc, anti-oxidants, e.g., carotenoids (such as a xanthophyll carotenoid like zeaxanthin or lutein), and micronutrients can be co-formulated or co-administered.

In addition, the aqueous composition according to the present invention is also useful for preparing a medicament such as a therapeutic or prophylactic agent for the VAP-1 associated diseases.

Examples of the preferred compound (I) thiazole derivative according to the present invention are listed in the following tables.











No.	Structure	No.	Structure
1		11	
2		12	
3		13	
4		14	
5		15	
6		16	
7		17	
8		18	
9		19	
10		20	











No.	Structure	No.	Structure
21		26	
22		27	
23		28	
24		29	
25		30	











No.	Structure	No.	Structure
31	 HCl	36	 HCl
32	 2HCl	37	 HCl
33	 HCl	38	 HCl
34	 HCl	39	 HCl
35	 HCl	40	 HCl











No.	Structure	No.	Structure
41	 <chem>CN1CCN(CC1)C(=O)N2C(=O)N(C)C(=O)N2C3=NC(=N)SC3C4=CC=CC=C4NC(=N)N</chem> HCl	46	 <chem>COc1ccc(cc1)Cc2c3c(ccn3S2)C(=O)N(C)C(=O)N2C(=O)N(C)C(=O)N2C3=NC(=N)SC3C4=CC=CC=C4NC(=N)N</chem> HCl
42	 <chem>CN(C)C1CCN(CC1)C(=O)N2C(=O)N(C)C(=O)N2C3=NC(=N)SC3C4=CC=CC=C4NC(=N)N</chem> HCl	47	 <chem>COc1ccc(cc1)Cc2c3c(ccn3S2)C(=O)N(C)C(=O)N2C(=O)N(C)C(=O)N2C3=NC(=N)SC3C4=CC=CC=C4NC(=N)N</chem> HCl
43	 <chem>CN1CCN(CC1)C(=O)N2C(=O)N(C)C(=O)N2C3=NC(=N)SC3C4=CC=CC=C4NC(=N)N</chem> HCl	48	 <chem>COc1ccc(cc1)Cc2c3c(ccn3S2)C(=O)N(C)C(=O)N2C(=O)N(C)C(=O)N2C3=NC(=N)SC3C4=CC=CC=C4NC(=N)N</chem> HCl
44	 <chem>CN1CCN(CC1)C(=O)N2C(=O)N(C)C(=O)N2C3=NC(=N)SC3C4=CC=CC=C4NC(=N)N</chem> HCl	49	 <chem>CCOC(=O)c1ccc(cc1)Cc2c3c(ccn3S2)C(=O)N(C)C(=O)N2C(=O)N(C)C(=O)N2C3=NC(=N)SC3C4=CC=CC=C4NC(=N)N</chem> HCl
45	 <chem>COc1ccc(cc1)Cc2c3c(ccn3S2)C(=O)N(C)C(=O)N2C(=O)N(C)C(=O)N2C3=NC(=N)SC3C4=CC=CC=C4NC(=N)N</chem> HCl	50	 <chem>COc1ccc(cc1)Cc2c3c(ccn3S2)C(=O)N(C)C(=O)N2C(=O)N(C)C(=O)N2C3=NC(=N)SC3C4=CC=CC=C4NC(=N)N</chem> HCl











No.	Structure	No.	Structure
51		56	
52		57	
53		58	
54		59	
55		60	











No.	Structure	No.	Structure
61	 HCl	66	 2HCl
62	 HCl	67	 2HCl
63	 HCl	68	 HCl
64	 2HCl	69	 HCl
65	 2HCl	70	 HCl

No.	Structure	No.	Structure
71	 HCl	76	 HCl
72	 HCl	77	 HCl
73	 HCl	78	 2HCl
74	 HCl	79	 2HCl
75	 HCl	80	 HCl



No.	Structure	No.	Structure
81	 HCl	86	 HCl
82	 HCl	87	 HCl
83	 HCl	88	 HCl
84	 2HCl	89	 HCl
85	 HCl	90	 HCl

No.	Structure	No.	Structure
91		96	
92		97	
93		98	
94		99	
95		100	

No.	Structure	No.	Structure
101	 2HCl	106	 HCl
102	 HCl	107	 HCl
103		108	 HCl
104		109	 HCl
105	 HCl	110	 2HCl

No.	Structure	No.	Structure
111	 <p>2HCl</p>	116	 <p>3HCl</p>
112	 <p>2HCl</p>	117	 <p>HCl</p>
113	 <p>2HCl</p>	118	 <p>HCl</p>
114	 <p>2HCl</p>	119	 <p>HCl</p>
115	 <p>2HCl</p>	120	 <p>2HCl</p>


No.	Structure	No.	Structure
121	 2HCl	126	 2HCl
122	 2HCl	127	 2HCl
123	 2HCl	128	 2HCl
124	 2HCl	129	 HCl
125	 2HCl	130	 2HCl

No.	Structure
131	
132	


The present invention is further illustrated by means of the examples shown below. The examples, however, should not be used for limiting the scope of the invention in any means.

Test Example 1


Compound A:



Compound B:



Compound C:



Compound A was added with distilled water (injection

grade) and stirred to give an aqueous saturated solution of Compound A (3.55 mg/mL, about pH 7) .

Compound B was added with distilled water (injection grade) and stirred with adding hydrochloric acid to give an aqueous saturated solution of Compound B (1.12 mg/mL, about pH 3) .

Compound C was added with distilled water (injectable grade) and stirred with adding hydrochloric acid to give an aqueous saturated solution of Compound C (10.73 mg/mL, about pH 7) .

Sodium chloride was added to thus obtained aqueous saturated solution of Compound A, B, and C respectively in an amount to give 0.4% NaCl solution and the mixture was stirred. Thus obtained mixture was observed for precipitation and the solubility, i.e. the concentration of the Compound A-C in the solution or supernatant (mg/mL) .

Similarly, the saturated aqueous solution of Compound A-C was added with sodium chloride in an amount to give 0.85% solution, and precipitation and solubility were observed.

Results are shown in table 1 below.

TABLE 1 effect of sodium chloride on the solubility of the Compound A, B and C

	conc. of NaCl (%)	Precipitation	Solubility (mg/mL)	% Solubility
A	0	no	3.55	=100
	0.4	yes	1.11	31
	0.85	yes	0.69	20
B	0	no	1.12	=100
	0.4	no	1.17	100
	0.85	no	1.19	100
C	0	no	10.73	=100
	0.4	yes	1.76	16
	0.85	yes	0.86	8

Test Example 2

To the saturated solution of Compound C prepared in the same manner as test example 1, glycerin in an amount to give 2.5% solution was added and the mixture was stirred. Similarly, mannitol in an amount to give 3.5% solution, boric acid in an amount to give 2% solution, potassium chloride in an amount to give 0.2% solution, disodium hydrogenphosphate in an amount to give 1% and sodium citrate in an amount to give 1% solution were added to the aqueous saturated solution of Compound C respectively. Precipitation and solubility, i.e. the change of the concentration of Compound C in the solution or supernatant were observed.

Table 2: Effect of the additives on solubility of Compound C.

additives, conc. in the solution (%)		precipitation	Conc. of C w/o additive (mg/mL) (A)	Conc. of C /w additive (mg/mL) (B)	B/A ratio (%)
Glycerin	2.5%	no	11.51	10.91	95
Mannitol	3.5%	no	11.51	11.05	96
Boric acid	2%	no	11.51	11.59	100
Potassium Chloride	0.2%	yes	10.73	3.16	29
disodium hydrogenphosphate	1%	yes	10.73	2.54	24
sodium citrate	1%	yes	10.73	0.48	4

Test Example 3

Compound C was added with glycerin and distilled water and stirred with adding hydrochloric acid to give an aqueous solution (Conc. of Compound C: 0.3%, conc. of glycerin: 2.5%, about pH 6). Similarly, aqueous solutions comprising Compound C and the additives shown in the table 3 were prepared.


Thus obtained solution was stored at 40°C in the LDPE container and the concentration of Compound C in the solution was observed over time. Result is shown Table 3 below:

Table 3. Effect of the additives on the stability of the aqueous solution comprising Compound C

Additives, conc. in the solution (%)		Change in Concentration of Compound C (vs the initial conc.) (%)			
		Initial	40°C		
			1 month	3 months	6 months
Glycerin	2.5%	=100	105.4	110.6	112.3
Mannitol	4.7%	=100	103.7	108.4	109.6
Boric acid borax	1.68% 0.018%	=100	104.9	106.2	108.2

Test Example 4

5 Compound D



Compound D was added with glycerin and distilled water and stirred to give clear aqueous solution (Conc. of Compound D: 0.3%, conc. of glycerin: 2.5%, about pH 6). Similarly, aqueous solutions comprising the same concentration of Compound D and the additives shown in table 4 were prepared.

Thus obtained solution was stored at 40°C in the LDPE container and concentration of Compound D in the solution was observed over time. Result is shown Table 4 below:

Table 4. Effect of the additives on the stability of the aqueous solution comprising Compound D

Additives, conc. in the solution (%)		Change in Concentration of Compound D (vs the initial conc.) (%)			
		initial	40°C		
			1 month	3 months	6 months
Glycerin	2.5%	=100	100.8	107.7	104.4
Boric acid borax	1.68% 0.018%	=100	99.7	105.8	102.6

INDUSTRIAL APPLICABILITY

5 The present invention provides an aqueous composition comprising a thiazole derivative of the formula (I): $R^1-NH-X-Y-Z$ (I)

wherein each symbol is as defined above, or a pharmaceutically acceptable salt thereof, and an additive
10 selected from the group consisting of polyol, sugar, sugar alcohol, boric acid or its salt, and water.

The aqueous composition of the present invention is very stable and therefore, aqueous dosage formulae which can be stably stored for a long time can be provided.